

Concentration

Percent concentration by mass $\%(m/m)$

$$\% \text{conc.} = \frac{\text{mass solute}}{\text{mass solution}} \times 100$$

Percent concentration $\%(m/v)$

$$\% \text{conc.} = \frac{\text{grams solute}}{\text{mL solution}} \times 100$$

Parts per million by mass

$$\text{ppm} = \frac{\text{mass solute}}{\text{mass solution}} \times 10^6$$

Percent concentration by volume $\%(v/v)$

$$\% \text{conc.} = \frac{\text{volume solute}}{\text{volume solution}} \times 100$$

For the concentration equations, mass and volume can be in any units as long as the numerator and denominator match.

Molarity (M)

$$M = \frac{\text{moles solute}}{\text{liters solution}}$$

Molality (m)

$$m = \frac{\text{moles solute}}{\text{kg solvent}}$$

Dilution

$$c_1V_1 = c_2V_2$$

Gas Solubility and Pressure (Henry's Law)

$$\frac{\text{solubility}_1}{\text{pressure}_1} = \frac{\text{solubility}_2}{\text{pressure}_2}$$

Colligative Properties

Freezing Point Depression

$$\Delta T_F = i \cdot K_F \cdot m$$

ΔT_F - Freezing point depression; i - # ion particles per solute molecule; m - molality; K_B - Cryoscopic constant of the solvent

Boiling Point Elevation

$$\Delta T_B = i \cdot K_B \cdot m$$

ΔT_B - Boiling point elevation; i - # ion particles per solute molecule; m - molality; K_B - Ebullioscopic constant of the solvent

Osmotic Pressure

Note that osmotic pressure uses molarity, not molality.

$$\Pi = iMRT$$

Π - osmotic pressure, atm; i - # ion particles per solute molecule; M - molarity; T - temperature, °K; R - ideal gas constant, 0.08206

K_F & K_B for water

$$K_F = 1.86 \text{ } ^\circ\text{C kg/mol} \quad K_B = 0.512 \text{ } ^\circ\text{C kg/mol}$$