

## Hypotheses

---

### Null vs Alternative Hypotheses

- **Null Hypothesis,  $H_0$** , is the claim regarding the population statistic that we are examining.
  - ▷ It is stated using the following notation
    - $H_0$ : characteristic = value
- **Alternative Hypothesis,  $H_a$** , is the competing claim; we are trying to find evidence for this.
  - ▷ It will have one of three forms:
    - $H_a$ : characteristic < value
    - $H_a$ : characteristic > value
    - $H_a$ : characteristic  $\neq$  value

### P-values and Proving $H_a$

- The **P-value** of a test result is the probability of that result occurring if  $H_0$  is true
  - ▷ We compare this to an agreed-upon threshold (the **significance level,  $\alpha$** ).

**P-value <  $\alpha$**  – we have convincing evidence that  $H_a$  is true.

**P-value  $\geq \alpha$**  – we have failed to disprove  $H_0$  and we cannot take  $H_a$  as true.

Note that we have *not* proven  $H_0$ ; we have simply been unable to disprove it.

### Type 1 & Type 2 Errors; Power

- **Type 1 Error** - Reject  $H_0$  when  $H_0$  is actually true
  - ▷ *i.e.*, falsely accept  $H_a$
  - The significance level,  $\alpha$ , is the probability of getting a type 1 error.
- **Type 2 Error** - Not rejecting  $H_0$  when  $H_0$  is actually false
  - ▷ *i.e.*, Rejecting  $H_a$  when  $H_a$  is actually true
  - $\beta$  is the probability of getting a type 2 error
- **Power** - The probability that a test will correctly confirm  $H_a$  when  $H_a$  is, in fact, true.
  - ▷ Power =  $1 - \beta$

## Testing a Population Proportion, $\pi$

### Calculating a P-value

- We calculate the **P-value** of a test result by calculating the z-score of our result, compared to the hypothesized value, and then finding the probability associated with this z-score.
- The **test statistic** (z-value) of our test result is:

$$z = \frac{\text{statistic} - \text{parameter}}{\text{stddev of statistic}} = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$$

$Z$  - Z score/test statistic;  $\hat{p}$  - proportion in sample  
 $p$  - hypothesized proportion;  $n$  - sample size

- $H_0$  should be rejected if the P-value  $\leq \alpha$ 
  - $\alpha$  is the significance level chosen for the test.

#### Validity requirements

This equation is valid if:

- Random data
- 10% rule  
 $n \leq 0.1N$
- Large counts  
 $n \cdot \hat{p} \geq 10$   
 $n(1 - \hat{p}) \geq 10$

### Obtaining P-value from z score

- Upper-tailed test**       $H_a: \pi > h$       P-value =  $1 - P(z)$
- Lower-tailed test**       $H_a: \pi < h$       P-value =  $P(z)$
- Two-tailed test**       $H_a: \pi \neq h$       if  $z > 0$ , P-value =  $2(1 - P(z))$   
if  $z < 0$ , P-value =  $2P(z)$

$\pi$  - Test value from  $H_0$ ;  $h$  - hypothesized value;  
 $z$  - Z score/test statistic;  $P(z)$  - P-value from z table

#### Calculator Note

On your graphing calculator, three functions are associated with testing hypotheses:

- 1-PropZTest**      Proportion test
- T-Test**      Mean ( $s_x$ )
- Z-Test**      Mean ( $\sigma$ )

## Testing a Population Mean, $\mu$

### Calculating the test statistic

The test statistic is a  $t$  score when testing a population mean.

$$t = \frac{\text{statistic} - \text{parameter}}{\text{stddev of statistic}} = \frac{\bar{x} - \mu_0}{\frac{s_x}{\sqrt{n}}}$$

$t$  - t-score/test statistic;  $\bar{x}$  - proportion in sample  
 $s_x$  - sample standard deviation;  $n$  - sample size

#### Validity requirements

This equation is valid if:

- Random data
- 10% rule  
 $n \leq 0.1N$
- Large counts  
 $n \geq 30$